



HAL
open science

Stromal vascular fraction for the treatment of the radiation-induced gastrointestinal syndrome

Christine Linard, Lydia Bensemmane, Claire Squiban, Noëlle Mathieu, Fabien Milliat

► To cite this version:

Christine Linard, Lydia Bensemmane, Claire Squiban, Noëlle Mathieu, Fabien Milliat. Stromal vascular fraction for the treatment of the radiation-induced gastrointestinal syndrome. CBRNE Research Innovation conference, May 2022, Lilles, France. irsn-03980949

HAL Id: irsn-03980949

<https://irsn.hal.science/irsn-03980949>

Submitted on 9 Feb 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Stromal vascular fraction for the treatment of the radiation-induced gastrointestinal syndrome

Christine Linard, Lydia Bensemmane, Claire Squiban, Noëlle Mathieu, Fabien Milliat

Institute of Radiological Protection and Nuclear Safety, Fontenay-aux-Roses, France

Christine.linard@irsn.fr

Accidental or intentional radiation exposures have serious health consequences for exposed individuals and can affect a large number of people. Large volume irradiation at high irradiation doses induces multiple tissue lesions. The gastro-intestinal tract is particularly sensitive to irradiation and lethality. At dose more than 10 Gy results in diarrhea, dehydration, sepsis and intestinal bleeding with mortality within 10 post-exposure. Radiation-induced gastrointestinal syndrome (GIS) results from direct cytotoxic effects on intestinal stem cells and crypt stroma impairing epithelial regeneration. Damaged intestinal epithelium significantly reduces the mucosal integrity and promotes systemic bacteria influx resulting in sepsis and death. Given the logistical hurdle and the urgency for treatment in large numbers of casualties, there is a tremendous need for effective therapeutic measures, even if implemented several days after radiation exposure. The stromal vascular fraction (SVF) derived from adipose tissue is an easily accessible source of cells with angiogenic, anti-inflammatory, immunomodulatory, and regenerative properties. We examined whether SVF restores the irradiated intestinal cells niche and mitigates the GIS. At the day of abdominal irradiation (18Gy) mice were injected in systemic with SVF, obtained by enzymatic digestion of adipose tissue. Seven days post-irradiation, SVF treatment limited weight loss and inhibited intestinal permeability [1]. When injected before 24 hours post-irradiation, SVF limited the mortality. SVF has an anti-inflammatory effect in the intestine by repressing pro-inflammatory cytokines, accelerating the maturation of monocyte able to generate anti-inflammatory macrophages. Immunohistological analyses of intestine showed that SVF treatment stimulated the regeneration of the epithelium by promoting numerous hyperproliferative zones. SVF restored the cell population in the intestinal stem cell compartment. The *ex-vivo* intestinal “organoid” model that mimics the clinical response confirmed that SVF treatment stimulated the intestinal stem cell compartment. With pleiotropic effects that contribute to limit radiation-induced lethality, SVF offers attractive prospects for the treatment of emergency GIS.

[1]. Bensemmane, C Squiban, C Demarquay, N Mathieu, M Benderitter, B Le Guen, F Milliat, C Linard, *Stem Cell Res Ther*, 2021,12:309.